

The best time to plant a tree was 20 years ago.
The second best time is now.....



Minnesota SWCD Tree Planting Handbook

Dear Tree Planter . . .

With headlines reporting the continuing deforestation of the tropical rain forest, one may ask the question: Are America's forests in danger of disappearing? Because people such as yourself practice reforestation, our forested acres are actually growing in size. About one-third of the United States, or 731 million acres is covered with trees. That's about 70 percent of the forest that existed when Columbus discovered America. Almost a third of this is set aside in permanent parks and wilderness areas.

Minnesotans' have planted an average of 12 million trees annually, enough trees to cover over 15,000 acres per year. Good land stewards are planting trees for many good reasons. The results of their efforts can be seen in reduced soil erosion, improved air and water quality, healthy forest industries, enhanced wildlife habitat and generally a more attractive surrounding for us to live in.

Aspen has become the most prominent tree in Minnesota's forests. After clear-cutting, aspen regenerates readily by sprouting from its root system or by drifting seeds onto disturbed sites. Most of the other major species in Minnesota need some help from tree planters to ensure that they make up a part of the new forest.

The following pages will help explain how to plant and care for a tree seedling. There is a section on the general characteristics and planting requirements of the tree and shrub species commonly planted for conservation purposes in Minnesota.

The professionals working in conservation throughout Minnesota "Thank You" for planting, nurturing and wisely using one of Minnesota's greatest treasures-its renewable trees.

Table of Contents

Answers to Frequently Asked Questions	4 - 5
Planting for Wildlife Habitat	5 - 8
Planting Shelterbelts and Field Windbreaks	9
Shelterbelt Design	10
Shelterbelt Layout	11
Livestock and Field Windbreaks	12
Planting Trees for Energy Conservation	13
Planting Trees for Wood Production	14
Growing Christmas Trees	15
Cost-Share Assistance Is Available	16
Trees of Special Concern	17
Planting Sketch	18
Worksheet to Estimate Landowner Share of Cost	19
Before You Plant	20
Preparing the Site for Planting	21
Tree Planting	22 – 23
How to Plant a Tree	24
Common Mistakes That will Kill Seedlings and # of Trees to be Planted per Acre	25
Containerized Stock	26
Tree Shelters	27
Tree Maintenance	28
Tree Planting Check List	29
Introduction To Tree Care	30
Planting Success	31
Water and Nutrient Needs	32 – 33
Weed Control	34 – 35
Animal Depredation	36
Tree Diseases	37
Insect Pest Management	38
Tree Pruning and Thinning	39
Acknowledgement / Information	40 - 41

Sponsored by:

Minnesota DNR, Division of Forestry; Minnesota Association of Soil & Water Conservation Districts; The Minnesota Forestry Association; Natural Resources Conservation Service; Minnesota Forest Stewardship Committee; and your Local Forestry Committee.

Answers to Frequently Asked Questions

When is the best time for weed control on a new planting? **Answer:** On old field or cropland sites, fall application of herbicides seems to be the most effective. Your plan will suggest the herbicide treatment best to control grass and weeds on your site.

What's wrong with my trees, they're all turning brown? **Answer:** Call your forester, resource advisor or whoever helped you set up the planting immediately. It could be the result of many things - trees were not handled properly and dried out during the planting; you may have gopher problems; the dirt was not packed well around each seedling after planting; the seedlings may have a disease.

I'm concerned about the environment and the use of herbicides in my tree planting. Is there an alternative? **Answer:** Herbicides recommended for controlling weeds are federally registered for that use and are considered by the EPA as safe, if used properly for that purpose. There are alternatives such as cultivation, mulching or fabric weed barriers. The important thing is to keep an area grass and weed free around each seedling for at least 3 feet, until the trees are freely growing.

My trees just arrived and I can't plant them until next week, what should I do with them?
Answer: Make arrangements for cold storage (40° to 50°F) or heel-in the seedlings. See page 22 for a detailed description of "heeling-in". Storing the trees for a few days in a cold garage or fruit cellar is acceptable, but any longer will result in the seedlings rotting. Do not store seedlings in tubs of water - they will die.

It's already March, I haven't done anything about setting up a tree planting project. Is it too late?
Answer: If you want a hassle free tree planting the answer is YES! You should work on the site preparation this year and plant next spring. You can still get it done this year, however, you must be willing to spend a lot of time looking for tree seedlings, working on site preparation and waiting your turn for planting equipment. It is best to wait a year.

Should I have a contract with the vendor that plants my trees? **Answer:** It is always sound business practice to require a contractual agreement when you hire someone to do work for you. It is the final word to resolve disputes surrounding completion of the project. It can reduce your liability for the other party's actions.

Can I plant all my seedlings in a small plot and move them later when they are bigger?
Answer: It is best to plant the new seedlings where you want them. Every time the trees are transplanted they experience transplant shock resulting in lost growth causing a lot of extra work.

What is the best way to plant a site - direct seeding of seeds or planting seedlings, transplants or container grown seedlings? **Answer:** Each situation is different and it is best to consult with a forester or other resource advisor to receive some advice.

The estimate I received for my planting project was too expensive. Is there some means to defray these expenses? **Answer:** Yes. The first way is to apply for one of several cost-share programs available for planting trees and shrubs. These programs can cover 50% to 65% of the project costs. This leaves the owner to pay the remainder, or provide the labor or materials as in-kind contribution. Many times a landowner can provide enough labor to the project to offset their share. Your plan preparer can outline this in your project budget.

The trees in the boxes have some gray stuff on them. Is that OK? **Answer:** You may just be seeing a harmless mold on the seedlings. If the box has a strong sour, fermenting odor, your trees may be starting to rot because they were not stored properly. Call your forester or resource advisor immediately for further assistance.

I've ordered the trees and contracted the tree planter, but I forgot to apply for the cost-share program. Is it too late? **Answer:** Maybe not. Call the local agency office administering the cost-share program. If you haven't started actual work on the project, you may still be able to apply. It is up to the agency to decide on your eligibility.

Can I mix seedlings in my tree planting? **Answer:** In most cases the answer is yes and in general, it creates a more naturalized planting. However when planting a site for the sole purpose of timber production, it is not recommended to mix species in the rows. Some planting designs have species planted in alternating rows, like white pine and red oak or walnut, where the pine help train the hardwoods into better quality trees.

Should I worry about the old box elders by my tree planting site? **Answer:** Yes. Box elders and elm are prolific seed producers. Your site preparation not only is good for the establishment of your small seedlings, but is also an excellent site for seeds to germinate. It is recommended to remove all elm and boxelder around your site if possible by cutting and chemical treatment. Dead trees can be left standing for wildlife den trees.

I planted trees a few years ago, now all I see in my field is a sea of grass. What happened to my trees? **Answer:** It is important to monitor your tree planting regularly, at least every couple weeks during the first year. If it looks like weeds are beginning to grow over the small trees, contact your forester or resource advisor for further assistance on weed control measures. Mowing will help keep the grass from falling on the trees and smothering them.

What kind of records do I need to keep for a cost-share program? **Answer:** It is a good idea to start a journal tracking everything you do on the project. Keep track of all time spent working on the tree planting, all bills for materials and tree purchases and receipts for all labor and/or contractors. You will need to present these to the agency administering the cost-share program to verify your costs. These records should be maintained for income tax purposes.

Planting for Wildlife Habitat

Every animal has a different group of plants to which it is attracted. Therefore, selections for your planting should be made on the basis of meeting some of the requirements for the widest variety of desired animals. Other desirable animals can be attracted to the area by providing nesting structures and food plots.


Always consider the total annual needs of wildlife. Don't be concerned if you can't fit all needs on your property. By investigating the cover on neighboring lands you may find several cover types that will compliment your planned project and, overall, provide for the seasonal needs of your desired animals.

Throughout the Upper Midwest it is vital to concentrate on providing food and cover plants. Recent research in established shelterbelts indicates that white spruce, Colorado spruce, cottonwood, boxelder, and Siberian elm offer the most benefits to the widest range of animals in open agricultural areas. These trees are especially beneficial for birds during the nesting season.

Where possible, plant approximately 25% trees and shrubs that provide mast (acorns and nuts) and/or fruit. Deciduous trees such as maple, oak, walnut, hickory, hackberry, crabapple, plum and hawthorn are excellent species. Conifers such as spruce, cedar and fir are better selections than pine. Shrubs should include nut and fruit bearing plants; however, some animals will have an interest in eating the buds of these plants as well.

For wildlife habitat, when in doubt plant:

- Spruce or fir rather than pine
- Cottonwood or hybrid poplar as tall deciduous trees
- A mixture of red and white oaks, including bur oak
- Fruit and nut bearing plants also edible by humans
- Native species rather than exotics or ornamental cultivars
- More warm season grass (prairie) than cool season grass (brome and bluegrass)



Planting a Year-long Food Supply for Wildlife

The following shrubs and small trees are just a few of the species that are recommended for a general purpose wildlife cover planting. These plant varieties can survive on a wide variety of Minnesota soils and will provide a year-long supply of food and cover for many birds and animals.

Highbush Cranberry: This shrub ranges in size from 3 to 9 feet tall. Its tri-lobed, maple-like leaves are attached to red stalks. Its brilliant red berries provide food for wildlife as well as providing a pleasing visual sight during the winter months.

Caragana: An upright growing tall shrub used for windbreaks, erosion control, and food and cover for wildlife. A hardy shrub, it can adapt to most sites, but prefers open sunny locations.

Wild Plum: Used for food and cover for wildlife. Considered a fast growing small tree growing to a height of 12' to 20'. It is quite hardy, does well in full or partial sunlight, and tolerates both dry and wet soils. Densely planted, it will form a thicket that is good for wildlife habitat.

Crabapple: Trees with white to deep pink flowers. Fruit ranges in size and color from small yellow apples to medium-sized red crabapples. Many species of birds and animals rely on the seasonal crop of apples for food, including the cedar waxwing, purple finch, robin, red fox and cottontail rabbit. Crabapples do best in full sun to partial shade and on good, well-drained soils.

Nanking Cherry: A very hardy, medium sized shrub that produces showy white flowers. On occasion, severe cold winters can damage flower buds above the snowline, reducing the abundance of fruit produced. The small attractive red fruit is tart and is favored by cedar waxwings, squirrels and chipmunks. Nanking cherry will grow in sandy soil but does best in heavier soils, in full sun.

Red-osier Dogwood: Native to Minnesota, this shrub has been widely planted for its red winter stems. Dogwoods prefer fertile, moist soil and full or partial sunlight. The fruits are white and ripen late in the summer but can persist into the early winter months. Evening grosbeaks, chipmunks and white-tailed deer are just a few of the many birds and animals attracted to the dogwood. This shrub will spread naturally by root suckers.

Cotoneaster: Attractive to catbirds, mockingbirds and purple finches. Pinkish flowers bloom in early spring and black fruits ripen in late summer and early fall. It is susceptible to oyster shell scale and fireblight.

Juneberry: Also known as serviceberry or shadberry, this shrub is quite variable from plant to plant and is difficult to identify. All juneberries have white flowers and black-blue fruit that matures in July. Songbirds, ruffed grouse, mourning doves, skunk, beaver, fox, bear and squirrels are but a few of the many animals that eat the fruit. Many other animals browse the twigs. Juneberries like well- drained soil and full sun light although they will tolerate partial shade.

Eastern Red Cedar: A small to medium-sized conifer tree from 40' – 50' feet tall. The cone of this tree is actually a berry-like fruit that birds will eat in large numbers. The foliage of red cedar provides excellent winter cover for a variety of wildlife species. The natural range for red cedar includes the southern one-third of Minnesota. Red Cedar is the host for cedar-apple rust. Be wary of planting near prized apple trees or commercial apple orchards.

Chokecherry: Native to Minnesota, this hardy shrub can attain the height of a small tree. It produces white flowers and tart, purplish fruit that can be used to make jellies and pies. Songbirds, ruffed and sharptail grouse, pheasant, raccoon, black bear, red fox, white-tailed deer, cottontail rabbit and gray squirrel feed on its fruit and twigs. It prefers rich and moist soils and will grow in full sun light or shade.

Hawthorn: A dense shrub or small tree, the hawthorn has long, sharp thorns on its stems and branches. It produces showy white, sometimes pink, flowers and small, yellow to red, and apple-like fruit. The fruit often remains on the plant all winter, providing food for numerous birds and animals. The hawthorn's dense crown offers excellent nesting cover for many kinds of songbirds. It is not particular as to soil, but likes sunny sites.

Establishing Wildlife Cover Plantings

As with any planting, weed control is essential. Tilling, mulching, or pre-treating with a herbicide is recommended around each seedling to reduce weed competition and to maximize shrub and tree survival. Cost-share funds to reimburse landowners for wildlife shrub and tree plantings are generally available.




Windbreaks benefit wildlife in many ways.

Planting Shelterbelts and Field Windbreaks

A well designed and maintained shelterbelt adds to the value of your home or farmstead. Contact your local SWCD or NRCS for technical assistance.


A Good Farmstead Shelterbelt Will...

- Moderate the effects of hot, dry summer winds
- Beautify the home and farmstead, making them attractive places to live, for work and recreation
- Reduce the effects of cold winter winds on humans and animals
- Help to attract songbirds and game birds
- Reduce the amount of noise and dust from adjacent roadways
- Prevent snow from drifting around buildings, roads and walkways
- Save tractor fuel and labor, by less snow plowing
- Reduce livestock feed costs, Cattle protected from winter winds use less feed to keep warm, Thus, winter weight gains require less feed
- Protect feedlots, gardens and orchards from harsh weather
- Aid in establishing lawn and landscape plantings



Shelterbelt Design

Cross-section of a 10-row shelterbelt



Standard Plan: The indicated spacing between rows is only a guide and shows minimum spacing's. Make spacing at least 4' to 5' wider than the width of equipment used to maintain the planting. The more conifers the better; 3 rows is a minimum suggestion.

TABLE I

If you only have room for:

7 rows
6 rows
5 rows
4 rows
3 rows

Use one of these combinations:

1, 2, 3, 4, 7, 8, 9 or 1, 2, 3, 5, 7, 8, 9
1, 2, 3, 7, 8, 9 or 1, 3, 5, 7, 8, 9
1, 3, 7, 8, 9 or 1, 2, 4, 7, 8
1, 2, 7, 8 or 1, 4, 7, 8
1, 7, 8 or 7, 8, 9

Your local SWCD office can help you design a windbreak or shelterbelt on your property.

Shelterbelt Layout

(For reference, rows are numbered from west to east. See the diagram on page 10.)

Rows 1 – 2:

Shrubs: Plants in these first rows are planted 4' to 5' apart within the row and 10' to 15' between the rows. Caragana, lilac or cotoneaster are good choices for these rows.

Rows 3 – 4:

Tall Shrub or Medium Tree: These seedlings are planted 5' to 8' apart in the row and 10' to 15' from the shrub row. Row four should be 10' to 15' from row three. Recommended species include American plum, common chokecherry or flowering crab.

Rows 5 – 6:

Tall Deciduous Tree: Row five should be spaced 16' to 20' behind row four. Row six should be spaced 16' to 20' behind row five. Trees should be planted 10' to 16' apart within the row. Deciduous trees suggested here include green ash, hackberry, hybrid poplar, soft maple and eastern cottonwood.

Rows 7 – 8


Tall Conifer (evergreen): Row seven should be spaced 16' to 20' behind row six, and row eight, 16' to 20' from row seven. Here again trees are planted 10' to 16' apart. Suggested species in these rows are Norway, Ponderosa and white pine.

Rows 9 – 10

Medium Conifer (evergreen): These last rows are both located 16' to 20' from the previous row and trees are planted 10' to 16' apart within the row. Black Hills spruce, Colorado spruce, Norway and white spruce, plus white cedar are good choices.

When planting windbreaks and shelterbelts, always consider where the resulting snowdrift will occur. Plantings should be placed far enough away so drifting snow does not adversely affect buildings and roadways.

Livestock and Field Windbreaks



An example of a 160 acre farm designed to take full advantage of windbreaks and other woody plantings. With a little imagination and careful planning, landowners can enhance both wildlife benefits and economic returns from their land.

Planting Trees for Energy Conservation

Strategic planting for energy conservation has received much attention because of the multiple benefits provided. Owners of buildings well-shaded and sheltered by trees can see direct savings on heating and cooling bills. And whenever energy use is decreased, less fuel is burned and less air pollution is created.

Just how beneficial trees are for conserving energy depends a great deal on how they are used. While the aim of a community-wide planting is to increase tree cover, the trees at each site should be located as optimally as possible.

"West is Best" - A building with well insulated walls and roof gains most of its unwanted summer heat through east and west facing windows. The most effective way to use trees to keep a building cooler in summer is to plant trees that **shade west facing windows**.

"Up and Over" - Cold wind leaking into a building is the biggest cause of winter heat loss. Tall, dense trees are ideally suited to direct wind up and over buildings. To achieve the greatest year-round energy savings, **create a windbreak** to the west and north.

"Let the Sun Shine In" - In the upper midwest, the owner of a typical fully air- conditioned home will spend about ten times more to heat than to cool the home. Since there is a significant amount of energy gained from the winter sun shining through south- facing windows, **avoid trees south of windows**.

"The More the Merrier" - Areas of extensive pavement and buildings become summer heat islands and windswept tundra in the winter. **Increase tree canopy cover** throughout the neighborhood to reduce both extremes.

Planting Trees for Wood Production

In forest situations where natural regeneration of desirable species is not adequate or completely lacking, trees must be planted to start or improve the composition of the future stand. Objectives for forest plantings can include one or more of the following:

1. Produce high quality timber and improve stand composition
2. Improve wildlife habitat
3. Reforest land not suited for agriculture
4. Reduce soil and wind erosion
5. Enhance esthetics

Forest plantations are a big investment usually only generating income for future generations. Therefore, it is important to develop a sound plan for planting the site, outlining all the necessary steps to ensure success. With the help of a forester or resource advisor, the landowner can identify their long-term forestry objectives and then fit the proper tree species to the site.

Depending on the objectives, different planting strategies may be used. Plantings designed for optimum fiber production will have trees planted in rows to permit efficient future intensive management. The first thinning will remove entire rows to provide more room for tree growth.

Forest plantings for timber production can be modified to include several of the other listed objectives. A variety of trees can be planted in a 'naturalized' mixture. Even though they are still planted in rows, these random mixtures become more attractive to wildlife and are more aesthetically pleasing as they mature, while still producing quality timber.

Planting plans for timber production usually suggest planting trees at spacing's ranging from 10' x 10' for hardwoods and 6' x 8' for conifers. Exact spacing should be based on the species selected and the planned uses of the harvested materials. Some cost-share programs may be available to assist with establishing forest tree plantations.

Growing Christmas Trees

Christmas trees may be grown as a primary crop or be removed as thinning from a tree planting established for timber. Popular trees grown in Minnesota are Scotch, Norway and white pine, Colorado and white spruce, and balsam and Fraser fir.

Today's Christmas tree farmer must produce a high quality tree to compete in the Christmas tree business. Careful management and shearing of the trees will produce a well-shaped, compact tree that can generate good income.

The ideal shape of a tree should resemble a cone, wide at the base and tapering uniformly to the top. A naturally grown tree usually has thin foliage, is spindly and has too much distance between the branches - a tree better left to grow to full size.

After planting, trees being grown for Christmas trees must be cared for more intensely than regular conservation plantings. Weeds must be controlled to keep them from choking out the small trees. Tall weeds grow up into the trees, shading the lower branches and causing the needles to drop.

Shearing begins when trees are 24" tall, generally two to three years after planting. Pine must be pruned when their new growth, or 'candles', have elongated, but before they harden off (become woody). Spruce and fir can be sheared at any time of the year. Once shearing is started it must be continued annually.

Tree planters interested in growing Christmas trees should consult with their forester for more information.



Before planting, check with your State Christmas tree association about the market for Christmas trees in your area.

Some markets are saturated.

Cost-Share Assistance is Available

Several government sponsored programs are available each year to help share the expenses of planting trees and shrubs for conservation purposes. Cost-share assistance rates range from 50% to 65% of the total eligible costs of tree and shrub planting projects. Most programs will allow in-kind contributions in the form of landowner- provided labor, equipment or supplies to reduce the amount of out-of-pocket expense for the project. See page 18 for a worksheet that will assist in calculating how much a landowner might pay for a planting project.

It is best to plan ahead at least one year for your tree planting project. This will provide the necessary time to meet with a forester or other conservation professional to analyze your site and prepare a plan. Cost-share programs generally have specific sign-up periods.

You must complete application for the cost-share program at the county USDA Service Center or an office of the local Soil and Water Conservation District. If you are relying on cost-share assistance for your planting, **do not begin the practice until you receive written approval from the office to which you've made application.**

After approval you can begin with the rest of the necessary tree planting project steps as outlined in your tree planting plan, such as:

- Ordering tree seedlings
- Make site preparation arrangements
- Make arrangements for tree planting
- Submit bills for project to agency
- Make arrangements for plantation release, as needed

Be sure to consult with your local forester, conservationist or other resource advisor to receive more information on any cost-share programs available for your tree planting project. Your resource advisor can help you understand the process of using the cost-share programs. See the back page of this handbook for a description of cost share programs available as of the date of this printing.

Trees of Special Concern

We are in the midst of a quiet environmental crisis. New foreign plants and animals are slowly infiltrating and changing our ecosystems. Many introduced species are more aggressive than native species and take over native habitats. They pose increasing threats to our lakes, parks, forest lands, wildlife areas, roadsides, agricultural lands and our nations native ecological balance. Human actions, intentional or accidental, are the main source of invasive species introductions. Potentially, any plant or animal species from places outside of Minnesota presents risks to our native ecosystems. The pet industry, bait dealers, recreational boating, biological supply houses, bird seed importation, grain shipments, horticultural introductions and international shipping can all bring exotic species to Minnesota. The growth of the international economy and increased trade in plants and animals have resulted in more and more accidental and deliberate releases of non-native species. Please think about the potential effects if you consider planting any of the below listed species.

Ginnala or Amur Maple

This is a shrub that readily reproduces by seed and has the potential to be invasive in some situations. It shades out prairie plants and if cut, will sprout from the stump.

Siberian Elm

This tree flowers early in spring before leaves unfold and the seed matures quickly. Seed is disseminated by wind, germinates readily and the seedlings grow rapidly. Siberian elm is an aggressive plant.

European Buckthorn


This is a shrub that is still sold as an ornamental but has naturalized and is found in old fields, fence rows and in forested understories. It readily reproduces by seed and is very invasive.

Norway Maple

This is a tree commonly planted for its varietal differences in foliage color both during the growing season and fall color season. It reproduces readily by seed and is aggressive.

Planting Sketch

Scale = _____



Worksheet to Estimate Landowner Share of Planting

Estimated site preparation cost:

Cost per acre: _____ # of acres _____ = \$ _____

Seedling Cost:

#1: Species: _____

Cost per seedling \$ _____ x # of seedlings _____ = \$ _____

#2: Species: _____

Cost per seedling \$ _____ x # of seedlings _____ = \$ _____

#3: Species: _____

Cost per seedling \$ _____ x # of seedlings _____ = \$ _____

Estimated Planting Cost:

Acres _____ x seedlings per acre _____ x \$ _____ per seedling = \$ _____

Shipping Cost of Seedlings = \$ _____

Total Estimated Project Cost: \$ _____

Estimated Cost Share:

Cost Share % _____

X

Estimated Project Cost _____

Estimated Contribution by Cost Share = \$ _____

(Do not exceed maximums available for each component)

Landowner "Out-Of-Pocket" Cost:

Total Cost \$ _____ - Cost Share \$ _____ = \$ _____

Note: completion of this worksheet does not guarantee cost share assistance. Individuals will need to complete an application for the program they are interested in.

Before You Plant

People have a general interest in wanting to plant trees. Each year millions of trees are planted for various purposes throughout Minnesota. Most farms have areas of marginally tillable land, odd hard-to-farm corners, old retired pastures and other similar areas. All of these provide an opportunity for tree planting.

In addition, any home or farmstead on an open exposed site has the need for a protective windbreak planting. The small area it takes for a properly designed windbreak will more than pay for itself in energy savings while providing much needed habitat for wildlife.

Once the decision is made to dedicate an area to planting trees, the area needs to be analyzed to determine the best species suited to growing on the site. **There is nothing more important than putting the right tree in the right place.** Not only does the soil type need to be identified, but other factors such as water table and past agricultural practices need to be identified.

When the tree species list has been narrowed down, the landowner must make the decision as to what the primary purpose of the planting will be. With this information in hand, a forester or other resource advisor can assemble a planting plan for the project, outlining step-by-step, the necessary procedures to be followed to assure a successful planting.

Order Trees Early

Every year a large number of landowners order tree and shrub seedlings in the fall or early winter. This means that the inventories in most nurseries becomes extremely limited by late winter and early spring. We urge you to order as soon as you have a planting plan to reserve your seedlings, even though you don't expect shipment until spring.

Many nurseries offer Early Bird Discounts!

Preparing the Site for Planting

Site preparation is one of the most important, yet most neglected aspects of tree planting. Site preparation helps eliminate weed and brush competition that can kill or greatly reduce growth of newly planted tree seedlings. As the name implies, it simply means getting the soil ready for tree planting. There are several methods that can be used for site preparation depending on the existing condition of the area.

Mechanical site preparation uses some sort of equipment to remove competing vegetation from the site. Examples include plowing a furrow with a tractor, using a bulldozer with a special blade or a brush cutter.

Chemical site preparation involves the application of a forestry-use labeled herbicide to competing vegetation. Chemicals can be sprayed in bands, in spots or broadcast over the entire site. Different chemicals are used in different situations. **Be sure to read and follow the label before application.**

A combination of chemical-mechanical site preparation may be necessary in some instances. This is usually the case when a site has been mechanically cleared of existing vegetation to expose bare mineral soil. This also applies when former agricultural fields are converted back to a tree cover. Until the site is planted to trees, a cover crop can be a useful practice to reduce weed pressure and prevent soil erosion. Herbicides are then used to control competing vegetation while the tree seedlings are becoming established.

Release is a very similar activity which may be incorporated after tree planting to control weeds. The *release* of small seedlings can be accomplished by either chemical or mechanical methods.

A good planting plan considers both site preparation and release methods prior to ordering the trees. Without considering control of competition, the new plantation may not survive and end up as a big expense.

Tree Planting

Choose Your Planting Method

Trees may be planted by hand or with a tree planting machine. Usually all plantings up to 1,000 trees are more efficiently planted by hand with a planting bar. Projects using over 1,000 seedlings should consider using a planting machine pulled by a tractor. In most counties, either DNR Forestry or the Soil and Water Conservation Districts have planting bars and machines available. It is a good idea to reserve this equipment well in advance of tree delivery.

Line up Your Planting Crew


Tree planting is something even the beginner can attempt. Usually about 500 trees can be planted per person per day with a planting bar. A tree planting machine can plant 4,000+ trees per day and needs a crew of two or more. When it comes time to plant, family members and friends often are willing to pitch-in. There's always something for everyone to do. Even with a two-man planting machine, it is suggested that someone walk behind and tamp in each tree.

If you are unable to tackle the entire project yourself, there are local vendors available for contracting part or all of the job. A list of vendor names is usually available from your forester or resource advisor.

Planting Steps:

1. Decide why you're planting trees and select those that fit your needs. Give careful consideration to the planting site. Picture how the trees will look when they mature and how much space each tree will occupy.
2. Planting tree seedlings in the spring provides the best results. Planting can be started as soon as the ground is free from frost and can go as long as the tree seedlings have not become moldy or broken dormancy.
3. **The time between taking tree seedlings from their nursery "beds" and planting them at a permanent site is very critical.** Anything that can be done to shorten this time period will **increase** survival chances of the tree seedlings. *This means keeping the packaged seedlings in a cool, dark location and planting them as soon as possible.*
4. At all cost, protect tree seedlings at the planting site! **Anything that causes tree seedlings to warm up or dry out before they are planted will cause damage to the seedlings.** Exposing roots to hot sunlight and drying winds for just three minutes can cause seedling mortality. Using a *planting gel* will protect the roots from drying while planting.
5. After receiving seedlings, all efforts should be made to plant them within 24 hours. If this is not possible, place the seedlings under refrigeration, at a temperature ranging from 40° to 50° F.

6. Be sure the seedlings are planted straight, with roots fully spread directly below the stem. It is very important to pack the earth down firmly. Check occasionally by pulling on a plant. Any air pockets can cause the tree to dry out and die.
7. Be sure to plant at the proper depth. Set the plant into the ground at the same level it formerly grew in the nursery. Never plant any deeper than the root collar found on the stem at ground level.
8. *Heeling-in seedlings.* If you need to postpone your planting operations for a week or more, it's best to heel-in the seedlings. This is done by digging a trench in a shady location and spreading out the seedlings. Fill the trench with dirt and water the seedlings well.




Be sure to “heel-in” bare root stock if planting is delayed for more than 48 hours.


How to Plant a Tree



1. Plunge planting bar or shovel into ground at the angle shown and push inward to upright position.



2. Insert tree at same depth it grew in nursery. If planted too deep, roots will curl; too shallow, roots will be exposed to the air and both errors will cause death to the seedling. Extra-long roots can be trimmed with a *sharp* knife to better fit the planting hole.



3. On correctly placed seedling, plunge bar or shovel into ground about 3" – 4" from hole.



4. Work bar forward and pull back. Using your feet, tamp and fill in last hole and any remaining loose soil around tree roots to prevent air pockets.

Common Mistakes That Will Kill Seedlings

- store seedlings in a bucket of water for more than a few minutes
- planting seedling too deep or too shallow
- allowing roots to curl back toward the top of the hole
- not allowing proper root spread
- planting in sod without good site preparation
- leaving trees in boxes exposed to the sun
- planting in dry soil
- planting a species not adaptable to the site
- planting where brush competition will crowd out the seedlings
- planting seedlings where they won't receive sufficient sunlight
- keeping trees in boxes more than a few days without cold storage

Number of Trees to be Planted Per Acre

Normal spacing for conifers result in 600-800 trees per acre. Hardwoods are planted at a density of 400-500 trees per acre. Shrub plantings are generally in rows 6' apart.

<u>Distance Apart</u>	<u># of Plants</u>	<u>Distance Apart</u>	<u># of Plants</u>
5 x 5 ft	1,742	10 x 10 ft	436
6 x 6 ft	1,210	10 x 12 ft	363
6 x 8 ft	908	12 x 12 ft	303
7 x 7 ft	889	Windbreak and Wildlife PlantingsCover	
7 x 8 ft	778	6 x 12 ft	605
8 x 8 ft	681	6 x 15 ft	484
8 x 10 ft	545	10 x 20 ft	218
9 x 10 ft	484	20 x 20 ft	109

Containerized Stock

Containerized tree seedlings are started in containers, usually under greenhouse conditions, and grown through one or more flushes before they are shipped to the field. They may be shipped in the container to the field or extracted as a plug and shipped. Containerized seedlings have some advantages over bare root planting stock:

1. The planting season can be extended into early fall because the root system of each seedling is protected by a plug of soil. This infers that the stock can also be held in storage for longer periods of time making scheduling of planting jobs that much easier.
2. Planting *shock* is minimal when trees are removed from the container and planted. The containers are an easy way to keep roots moist and trees from drying out.
3. Seedlings at all stages of growth may be shipped and planted from the container (dormant, hardened-off or actively growing).
4. The system is nearly fool-proof in that the seedlings are buffered against poor handling and storage by the container. Being a plug also avoids some planting problems such as “J-rooting” or splitting roots at the base of the crown. Because of these inherent features of the system, survival of containerized planting stock usually runs very high.
5. Seed sources and species identification are easily maintained using containerized seedlings. Seedlings are not removed from their container until planting and the container can easily be labeled to keep sources and species information available right to the planting site.

Containerized stock is easy to plant. It can be planted with the same tools used for bare root planting, such as planting bars, hoe-dads, planting dibbles, shovels or tube planters, as well as by machines. Consult a resource advisor to match the correct tool to your particular site. All planting stock should be kept moist and planted to the correct depth. The root plug should be firmed in and the top of the plug covered with native soil. This is an important step because the peat plug can act as a wick and dry the seedling out if left exposed. By covering the top with a thin layer of native soil, the plug attracts water like a sponge and becomes a water reserve for the tree seedling.

Tree Shelters

Tree shelters are a new technology being used in conservation plantings to improve survival and increase growth of young seedlings. Several manufacturers are currently marketing tree shelters.

A tree shelter is a plastic polymer tube, approximately 3½" diameter. The tubes are sold in various heights for use in a variety of planting situations. Each tube must be supported by a rot-resistant stake. A nylon mesh sleeve is placed over the top to prevent birds from becoming trapped in the tube. The tubes should be left on the tree. Sunlight will degrade them after several years and not affect the growing tree.

The tree shelter's purpose is to provide protection from damage and to maintain an ideal growing environment until the young seedling grows out of the top of the tube. The shelter protects the seedling from animal, chemical and mechanical damages. It traps moisture from overnight higher humidity and keeps the young seedlings leaves cool and moist. This protection from damage and reduced moisture stress results in accelerated growth.

How many tree shelters should be used in a planting? In a typical forest regeneration planting, the most common recommendation is to use 50-150 tree shelters on the 435-680 trees planted per acre. Not every tree planted will have a shelter. The goal is to put shelters on trees that are hard to grow or those species that will become the future high quality, high value trees in the stand. The trees to be helped by tree shelters could be planted at 18-30 foot spacing. The remaining areas on the site can be filled with trees that are less valuable but easier to grow.

Cost-Share Available for Tree Shelters

Planting projects using tree shelters are eligible for cost-share funds.

Funds are limited to 150 shelters per acre.

Ask your forester for more information.

Tree Maintenance

Successful tree growers know that weeds and grass steal growth from trees. In some regions of Minnesota, annual precipitation can barely support tree growth. Unwanted vegetation competes with trees for soil moisture, nutrients and growing space. Where weed competition is severe, a high percentage of trees die in the first few years. Many surviving trees may be stunted and deformed. Several methods are available to combat weed competition and give young seedlings a chance. Most methods are eligible for cost-share assistance.

1. **Chemical Weed Control:** Herbicides are often the most effective and economical method for controlling weeds in a large planting. By applying the proper herbicide, weeds are controlled during the active growing season, giving the seedling all the available moisture and nutrients. Herbicides can be applied to the entire planting site, in bands over the tree rows or to 4' diameter spots around each seedling.
2. **Tree Mats:** These special fabric weed barriers are made from tightly woven material that blocks weeds from growing around young seedlings. Tree mats have been shown to be effective for up to three years. Several manufacturers sell pre-cut 3' x 3' squares. Weed barrier fabric can also be purchased in bulk rolls from local landscape nurseries. Mats need to be fastened to the ground to be effective.
3. **Mulch:** On smaller plantings, placing a 3' - 4' circular area of mulch around each seedling will reduce weed competition and increase soil moisture. Wood chips, *ground* corn cobs or other coarse natural materials can be used. Do not use grass clippings or straw because they will attract rodents.
4. **Cultivation:** On old field plantings not prone to erosion, cultivation is an option. Because cultivation will promote germination of annual weeds, it must be continued throughout the growing season.
5. **Mowing:** Mowing tall weeds should be considered as a last option. Mowing does not limit the competition for water and nutrients by weeds. Weeds should be clipped if they have overgrown the young trees. Heavy winter snows will push the tall weeds and grass down and bury the small seedlings.

Tree Planting Check List

- Select a site
- Request information from a forester or other land manager
- Apply for a cost-share program, if applicable
- Choose a tree or shrub species suited to the site, with the assistance of a land manager or forester
- Order trees
- Choose and complete best method of site preparation
- Make arrangements for tree planting and plant trees as soon as they arrive in the spring
- Monitor tree growth throughout the summer
- Choose and complete fall release practice as needed

All of these activities should be done in consultation with a professional forester or other resource advisor to assure the greatest possible chance of success.

Introduction To Tree Care

The planting plan has been done, tree species have been selected, site prep has been finished, cost-share has been dealt with, and the tree planting project has been completed. I'm done now, right? 😊 Not exactly. Now that the tree plantation has been "born", the "care and feeding" phase has begun! This is a really fun time in the establishment of a young forest. The trees are planted and want to grow. You get to walk through occasionally and see how things are going!

The next section is intended to give you some pointers and direction on how your plantation should be growing and what you can do to help. The tree care section will cover how to measure planting success, water and nutrient needs, weed control, animal depredation, tree diseases, insect pest management, and tree pruning and thinning.

It's important to monitor and protect your investment. It's tough to be a seedling. Animals eat and strip off bark, branches, and roots. Insects like to consume most or all of the foliage on little seedlings. Diseases are waiting to do their thing on plantations, large and small. Weeds and other competing vegetation are robbing nutrients and moisture that would maximize seedling growth. And sometimes the weather doesn't want to cooperate either.

No matter how carefully things have been planned and carried out, it's still important to recognize that seedlings have undergone stresses in the process of moving from a nursery to their new surroundings under your care. With a little luck from Mother Nature and proper technique in handling and planting, those stresses were minimized. Your job is to monitor your plantation and keep an eye on how well the tree seedlings are adjusting and help counteract any problems they may be having. The following section should help get you started in the process known as plantation survival and follow-up.

Planting Success

The first step after planting trees is to assess the planting success. This involves determining what survived and what did not. You should check the planting immediately after the job is done to make sure all trees are firmly in the ground with good soil-to-root contact (no air gaps).

Why should trees be checked? 1) Ensure the survival of the trees planted. 2) Protect the investment of time and labor expended in the planting endeavor. 3) Ensure the planting plan will be accomplished.

When should trees be checked? As stated earlier, the planting should be checked right after completion. Then depending on the size of the planting, the purpose of the planting, and the number of trees planted. They should be checked at a minimum of once each year for the first five years. If the planting was done for wind or snow protection where it is important that each tree is in place, it is best to check more frequently. After the initial five years, the checks can be done every five years.

How should trees be checked? This will depend a great deal on the size of the planting. If it is a fairly small planting each tree can be checked. In large acreage forest plantings a random sampling should be done. A number of sample plots should be taken. Make a measuring stick that is 11.8' long. An 11.8' radius of a circle will give readings on 1/100 of an acre. A very simple method is to choose a compass heading and walk a few paces. Turn and face the direction that you came from and throw your hat back over your shoulder. This is done so you cannot pick a spot that you would like to sample. Then, using the hat as the center of the plot measure with the stick and check to see how many live seedlings are in the sample plot. The size of the planting will help determine the number of sample plots that should be taken. A usual procedure is to measure one plot for each acre of the tree planting. Take the total number of trees, divided by the number of plots taken, and then multiply by 100 to find the average number of live trees per acre estimated in the planting.

Normal mortality? Here again the size and the purpose of the planting need to be taken into consideration. In a large acreage forest tree planting it may be acceptable for 2 – 3 percent mortality the first year. Over the first five years 5 – 10 percent may be acceptable. On the other hand in a windbreak or living snow fence every tree is very important, so replanting is necessary if mortality occurs.

Recording / Summarizing Data:

It is a good idea to keep records of the information that you have accumulated on the tree planting. The information will be useful in the future.

New Tree Planting



Water and Nutrient Needs

WATERING

Trees require water, but improper watering can also cause serious damage to a tree planting. If possible, water as needed for the first few years after establishment.

Check your trees every week for the first growing season. The soil under the mulch should be moist but not soggy.

Here are some tree watering do's and don'ts:

1. Over watering forces oxygen out of the soil and results in oxygen starved roots. Newly transplanted trees need extra watering until their roots grow into the area beyond the planting hole. In general an inch of rainfall per week is adequate.
2. Water less often in late summer. Too much water late in the growing season may stimulate growth that will not have time to go dormant.
3. Avoid frequent light watering. This promotes the development of a shallow root system, which makes the tree susceptible to winter injury and summer heat stress.
4. Water with an open-end hose at a rate low enough to keep water from running off. A good rule is to water one hour per inch diameter of the tree. Tree roots extend away from the tree at least as far as the tree is tall, so place the hose at or beyond the drip line for most effective watering

FERTILIZING

Fertilization is probably not necessary if the growth rate and foliage color is normal for a particular variety. Newly transplanted trees may exhibit slow growth until they get re-established. Regular fertilization will help tree establishment in less than ideal planting sites such as very sandy or heavy clay soils. Fertilizing is also recommended for tree plantings that have suffered damage from insects or disease.

The best time to apply fertilizer is early April before new growth expands. However, fertilizers can be applied any time up to about mid- July. Application past that time may stimulate plant growth that will not have time to harden off before cold weather. An exception is the use of slow-release fertilizers. Application of slow-release fertilizers is usually safe from mid-summer through late fall. Never fertilize drought stressed trees. If conditions become dry after fertilizer applications it is very important to water the trees.



Fertilize with a complete fertilizer such as 10-8-6. This formula can vary somewhat, but usually the nitrogen content (the first number) will be higher than the phosphorous (second number) or the potassium (last number). Be aware of local restrictions on phosphorus applications. It is best to complete a soil test before fertilizing. Soil sample bags and directions are available at most county extension offices or check with the Soil and Water Conservation District. A soil test will also indicate whether the soil is acidic or alkaline. Some nutrients may be unavailable to the tree when the soil is too alkaline. Discuss fertilizer application and soil test recommendations with a forestry professional before application.

Fertilizer spikes are a convenient way to fertilize a small tree planting. However, since each spike contains a small amount of fertilizer, they are not cost-effective for larger plantings compared to granular products.

Weed Control

The most serious problem young tree seedlings face is competition from other plants and grasses. Some work at this stage can prevent stunted trees with slow growth rates. Unwanted vegetation competes with newly planted trees for moisture, nutrients, and growing space. Where weed competition is severe, a high percentage of trees will die the first few years. Site preparation before planting is the best way to control weeds, but several methods are available for weed control after the trees are planted. Most are eligible for cost-share assistance in the tree planting practice.

MECHANICAL

Tilling

This activity either destroys the weeds or limits its food producing capability. Tilling depth should be no greater than two or three inches deep and be outside the drip line of the tree. Tilling deeper than 3 inches will damage tree roots and hinder growth. Several passes each year may be required to control weeds.

Mowing

Mowing should be used in conjunction with other methods of weed control such as mulching. Mowing does not help increase moisture for trees, but it can reduce future competition. To provide effective control, plantings must be mowed before seed heads mature. If weed whip trimmers are used take caution not to allow contact with tree bark. Plastic tree guards protect the tree from accidental girdling and wounding.

Mulching

Mulching is an effective way of providing weed control. Mulches conserve moisture and can moderate soil temperatures. Organic mulches include straw, wood chips, and grass clippings. Sawdust, grass clippings, and other fine sized mulches can prevent air and water exchange and should be used with caution.

Place mulches 2” to 4“ deep and add more as the it decompose. Keep the mulch about 6” away from the base of the tree. This will discourage mice and other small rodents from chewing on the stem under the mulch.

Fabric

Typically called landscape fabric, this material comes in 3’ by 3’ squares or rolls. The fabric suppresses weeds while allowing for water and oxygen exchange. The edges of the fabric must be stapled for buried to hold it in place.



COVER CROP

Cover crops planted between tree rows can reduce the spread of aggressive weeds, allow for better rain infiltration, and provide organic mulch. If the cover crop begins to compete with the trees it must be harvested.

CHEMICAL

Herbicides can be used to control weeds in tree plantings. Selection will vary based on target weed type, growth stage and specie of tree. Some herbicides can be applied directly over the trees and others must be applied as a directed spray to minimize or avoid contact with trees. Special tools may be required for application such as sprayers, granular spreaders, and sponge applicators. Items to consider in herbicide weed control include soil characteristics, herbicide characteristics, depth to water table, and use patterns. Specific application for each herbicide is contained on the product label. Be sure to read and follow all label directions and consider environmental factors when choosing this very effective weed control method.

For more specific information about weed control visit:

www.extension.umn.edu/distribution/naturalresources/m1156.html

Animal Depredation

The number one animal responsible for feeding on newly planted trees in Minnesota is white tail deer. Not only do they browse on tree seedlings but as the trees get bigger damage from antler rubs can be detrimental to small trees. While there are several ways to reduce deer damage it is almost impossible to completely eliminate it.

Electric fencing can be effective but is expensive and difficult to maintain. Typically the fence is 7 strands and about 7' tall.

Tree shelters come in a solid tube or a mesh material and are held in place by a stake. While these options work well, they are often damaged by rubbing bucks, bears, and may require regular maintenance.

Additionally, tree growth inside the tubes is more easily damaged by early winter freezes and exceptionally warm early springs. Even so, tree tubes provide good protection from browsing deer. A homemade alternative is an 8" diameter 4' – 6' tall tube made from fence with 1" by 2" openings.

Repellants come in two forms, smell and taste. Both are available from lawn and garden stores and are effective. Because they deteriorate with time, repellants must be reapplied often.



Bud capping is a method of covering up the tempting tasty buds so the deer cannot eat them. Staple a small piece of paper over the terminal (top) bud as shown below. Although labor intensive, this method is very inexpensive.



Rabbits, voles, and mice can also be detrimental to trees. They chew the succulent young bark within reach all the way around the tree. The layer just below the bark called the cambium layer is where nutrients are transported between the leaves and roots. With this girdling of the bark the nutrient transport is cut off and the tree will die. One way to prevent this animal damage is with a wire barrier. Use a piece of hardware cloth at least 5' long by 2' wide. Put it around the tree to form a cage 2' tall and 1' in diameter. Set the cage slightly below the ground line and secure to a post.

Livestock can be very detrimental to a new tree planting and should be fenced out whenever possible. Other animals such as beaver, moose, and even squirrels can present a challenge to tree establishment. Talk with your forestry professional if you expect a problem with these or other animals.

Tree Diseases

There are many diseases a tree can develop. Disease problems are often complex and need professional assessment in order for the solutions to be successful. The best advice for this problem is to monitor your planting closely. At the first sign of problems such as dead or discolored foliage, or disfigured, abnormal stems or branches, contact the forestry professional that assisted in the planning of the planting and arrange for a field visit. The forester can look at the trees firsthand and do any sampling required to identify the problem. Solutions can then be discussed and implemented once proper identification has been made.

There are several good website resources for disease and insect identification (at the time of this printing).

www.bugwood.org This site has images of many tree pests and diseases and can get you started in identifying a problem in your stand.

www.dnr.state.mn.us/fid/ This is the Forest Insect and Disease Newsletter published by the Minnesota Department of Natural Resources.

www.extension.umn.edu/projects/yardandgarden/diagnostics/ This is an interactive web-based diagnostic tool for various plant diseases.



Insect Pest Management

There are almost as many insect pests as tree diseases. Closely monitoring your planting for early signs of problems is the best defense against wide spread insect damage. Signs of insects are similar to tree diseases. Defoliation and abnormal stem or branch growth are signs of problems. Insect pests are sometimes easier to identify than diseases because inspections of the tree planting can reveal specimens for identification. Again, the forestry professional is the best place to turn for assistance with this problem. Once identified, the pest control method can be developed to meet the needs of the planting.

The same resources used for tree disease identification may have information on insect identification.



Forest Tent Caterpillars

Tree Pruning and Thinning

Pruning and thinning are tools used to control diseases, develop strong tree structure, and assist a tree in getting rid of unproductive branches.

A healthy stand of trees requires occasional removal of diseased or insect infected wood (see Disease and Insect Management sections to identify infected wood). That may mean part of a tree or in some cases the whole tree needs to be removed. Branches removed should be disposed of properly to prevent reinfection.

If a tree is damaged such as broken branches, pruning those branches helps the tree to heal the wound left by the damage.


All woody plants lose branches as a response to competition. Branches that do not produce enough, through photosynthesis, to sustain themselves, die and eventually shed. Pruning can replace this natural process and improve the strength and longevity of the tree.

In general pruning should be done during the dormant season, usually the late fall to early spring before the buds begin to swell. Dead branches can be removed any time of the year. At planting, remove only diseased, dead, or broken branches. Shearing of evergreens should be done on new green growth, before the tree starts woody elongation. Prune young trees lightly. More mature trees can be pruned more heavily. Leave one-half to two-thirds of the canopy in all cases.

Prune to shape young trees, but don't cut back the leader. Remove crossing branches, branches that grow back toward the center of the tree, rubbing branches, branch stubs, suckers, and narrow weak crotches. As young trees grow, remove lower branches gradually to raise the crown, and remove branches that are too closely spaced on the trunk. Remove multiple leaders on evergreens and other trees where a single leader is desirable (timber production, shade trees).

For more information refer to US Department of Agriculture, Forest Service Publication NA-FR-01-95.

Thinning can be an effective tool in insect and disease control and for improving stand vigor. Consult a forestry professional for specific thinning recommendations for your tree planting.



Proper pruning cuts are made at the point where a branch or twig attaches to another. Cuts should be made so that only branch tissue is removed and the stem tissue is not disturbed.

Branches with strong U-shaped angles of attachment should be retained. V-shaped angle attachments should be removed.

For Additional Information Contact

The Minnesota Department of Natural Resources (DNR), Division of Forestry provides resource information, technical planning assistance, funding programs and general forestry administration.

Soil and Water Conservation Districts (SWCDs) help the public locate resource information and programs, technical assistance for woodland owners and funding for various woodland management activities.

The USDA Natural Resources Conservation Service (NRCS) offers cost-sharing for a variety of conservation practices including windbreaks, shelterbelts, living snow fences, tree planting and timber stand improvement.

The Minnesota Extension Service offers practical, research-based information on a wide range of forestry, forest products and wildlife management subjects. It provides information and formal education to the general public.

Acknowledgements

- ☒ MN Soil and Water Conservation District Forestry Association
- ☒ MN Board of Water and Soil Resources
- ☒ MN Department of Natural Resources
- ☒ MN Natural Resources Conservation Service
- ☒ University of MN Extension Service
- ☒ United States Department of Agriculture
- ☒ Forest Service
- ☒ SWCD's: Aitkin, Beltrami, Carlton, Crow Wing, Koochiching

This Printing Sponsored by:

MN Soil and Water Conservation District Forestry Association

Some cost share opportunities for private landowners:

Stewardship Incentive Program (SIP)

SIP provides cost share assistance so landowners can implement projects from site preparation to timber stand improvement activities. Apply to Farm Service Agency (FSA).

Forestry Incentive Program (FIP)

FIP is a cost share program intended to promote timber production. Only selected counties can participate in FIP. Apply to the Natural Resources Conservation Service (NRCS).

Minnesota Forestry Association (MFA)

Funds for MFA come from three sources; 1) donations from private individuals, forest industry and other groups; 2) the Legislative Commission on Minnesota Resources (LCMR); 3) the Minnesota general fund. This is a cost share program that can do many of the same practices done under SIP. Apply to DNR-Forestry.

White Pine Program (WP)

This program funds activities directly related to establishing white pine and enhancing white pine survival. Eligible items include planting, thinning, pruning, release, bud capping and any other activity which promotes white pine health and growth. Apply to DNR-Forestry.

Wildlife Habitat Improvement Program (WHIP)

This program focuses on wildlife enhancement and improvement. Apply to the Natural Resources Conservation Service (NRCS).

This publication was printed by McGregor Printing and Graphics in McGregor, MN.
All the paper used in this publication is FSC Certified.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, or marital or family status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.